arXiv:1611.06007v1 [hep-ex] 18 Nov 2016

November 21, 2016

Searches for top-antitop quark resonances at $\sqrt{s} = 13$ TeV with the CMS detector

Marino Missiroli, on behalf of the CMS Collaboration

Departamento de Física Teórica, Universidad Autónoma de Madrid
Ciudad Universitaria de Cantoblanco, 28049 Madrid, Spain

We present the first results on searches for new massive resonances decaying to a top-antitop quark ($t\bar{t}$) pair with the CMS detector at the LHC in proton-proton collisions with a center of mass energy of $\sqrt{s} = 13$ TeV. The data set considered corresponds to an integrated luminosity of 2.6 fb^{-1} recorded during the the first year of the LHC Run-2. Searches are performed by measuring the invariant mass distribution of the $t\bar{t}$ system in semileptonic and fully-hadronic final states. Dedicated techniques are used to identify the decay of highly-boosted top quarks in order to maximize the analyses' sensitivity for $t\bar{t}$ resonances with a mass above the TeV scale. No significant excess is observed in the data compared to the expected SM background and exclusion limits are set on the cross section of a $t\bar{t}$ resonance in various new physics scenarios.

PRESENTED AT

9^{th} International Workshop on Top Quark Physics
Olomouc, Czech Republic, September 19–23, 2016

The existence of new massive particles coupling to the top quark is predicted by numerous extensions of the Standard Model (SM). Examples of such theories include two-doublet Higgs models, theories predicting additional gauge interactions, e.g. models based on technicolor, and models with additional space-time dimensions. These new heavy particles could be observed at the Large Hadron Collider (LHC) at CERN as resonances in the invariant mass spectrum of a $t\bar{t}$ pair. In this document we report on the first results of two independent searches for $t\bar{t}$ resonances in pp collisions at a center of mass energy of $\sqrt{s} = 13$ TeV with data collected by the CMS experiment [1] at the LHC for a total integrated luminosity of 2.6 fb^{-1}. One search is performed on events with one muon or electron in the final state (semileptonic or lepton-plus-jets analysis) [2], while the other targets events in which both top quarks decay hadronically (all-hadronic analysis) [3].

Both searches are optimized to probe the existence of new $t\bar{t}$ resonances with masses well above the TeV scale. This requires the use of dedicated techniques for the efficient reconstruction of highly-boosted top quarks. In the case of the top quark hadronic decay, this corresponds to using a method to tag large-radius jets comprising all the decay products of the top quark decay and characterized by specific substructure properties. The two analyses make use of the same jet t-tagging algorithm, based on two jet variables: the groomed jet mass determined by the softdrop algorithm, shown in Figure 1, and the N-subjettiness ratio $\tau_{32} \equiv \tau_3/\tau_2$ [4]. The t-tagging selection is tuned to yield a t-mistag rate of approximately 3%.

The semileptonic analysis [2] selects events containing exactly one muon or electron, at least two high-p_T jets and missing transverse energy. The lepton isolation requirement is replaced by a selection based on the transverse momentum of the lepton with respect to the axis of its nearest jet. This method is found to be the most effective one to reject non-prompt and fake leptons from multijet production and preserve the efficiency for leptons coming from high-p_T top quarks. The full kinematic reconstruction of the $t\bar{t}$ system is performed using the reconstructed objects in the final state. For each event the best hypothesis is selected based on a χ^2 discriminator and an upper cut on the χ^2 value is finally applied to reduce the contribution of non-$t\bar{t}$ backgrounds. The main backgrounds in the final event sample are given by W + jets and $t\bar{t}$ production. Both backgrounds are modeled with MC simulation and their central predictions are corrected using data in dedicated control regions. The final event sample is split in exclusive categories based on the number of b-tagged and t-tagged jets in the event. Figure 2 shows the $M_{t\bar{t}}$ distribution in the two event categories with the highest signal efficiency. Good agreement between data and background expectation is found in the final $M_{t\bar{t}}$ spectra and no significant excess consistent with the presence of resonant $t\bar{t}$ production is observed.

The all-hadronic analysis [3] selects events with a dijet topology, requiring two large-radius high-p_T jets both passing the t-tagging selection. The main backgrounds in the final event sample come from QCD multijet production and $t\bar{t}$ production. The



Figure 1: Distribution of the softdrop mass of large-radius jets with $p_T > 500$ GeV, $|\eta| < 2.4$ and $\tau_{32} < 0.69$ for data and simulation in the semileptonic analysis [2]. The simulation prediction is shown separately for jets matched (red) and not matched (blue) to a top quark at generator level. The simulation error band includes only the MC statistical uncertainty.

latter background is modeled with a next-to-leading order MC simulation, while the former one is determined with a data-driven method based on the measurement of the t-mistag rate in data control regions. The $M_{t\bar{t}}$ distributions are measured in six exclusive categories with different signal sensitivities and background composition. The categorization is based on the value of the rapidity difference between the two jets (Δy_{jj}) and the number of b-tagged subjets in the events. The $M_{t\bar{t}}$ distributions in the two most sensitive categories, given by events with $|\Delta y_{jj}| < 1.0$ and at least one b-tagged subjet, are shown in Figure 3. No significant deviation from the expected SM background is observed in the mass spectra of the reconstructed $t\bar{t}$ pair.

Since no evidence of resonant $t\bar{t}$ production is observed, the $M_{t\bar{t}}$ spectra are used to set upper exclusion limits on the cross section of a $t\bar{t}$ resonance with a mass up to 4 TeV in various new physics scenarios. These include models with a $Z' \to t\bar{t}$ resonance with different width hypotheses and a Kaluza-Klein gluon resonance in extradimensional models. Figure 4 shows the cross section limits set by the two searches for one of these signal models. These upper cross section limits are also recast in terms of lower limits on the mass of a $t\bar{t}$ resonance. This is shown in Table 1, which also includes for comparison the corresponding limits set by the combination of $t\bar{t}$ searches performed by the CMS experiment at $\sqrt{s} = 8$ TeV [5]. Despite the limited amount of integrated luminosity, the lower mass limits set by these first Run-2 analyses are already competitive or even improved with respect to previous searches thanks to the increased center of mass energy of $\sqrt{s} = 13$ TeV reached by the LHC. In particular, the lower mass limits for large-width signals, e. g. Z' resonances with a width of 10%, are significantly extended.



Figure 2: Post-fit distributions of the invariant mass of the reconstructed $t\bar{t}$ pair in the semileptonic analysis for events without a t-tagged jet and with at least one b-tagged jet (left) and for events with a t-tagged jet (right). The signal template shown in each plot is normalized to a cross section of 1 pb. The uncertainties associated to the background expectation include statistical and systematic uncertainties. The ratio of data to SM background is shown in the bottom panel of each plot, where the statistical (light gray) and total (dark gray) uncertainties are displayed separately [2].



Figure 3: Post-fit distributions of the invariant mass of the reconstructed $t\bar{t}$ pair in the all-hadronic analysis, for events with $|\Delta y_{jj}| < 1.0$ and exactly one b-tagged subjet (left) or at least two b-tagged subjets (right). Each signal template is normalized to a cross section of 1 pb. The uncertainties associated to the background expectation include statistical and systematic uncertainties. The ratio of data to SM background is shown in the bottom panel of each plot [3].



Figure 4: Expected and observed 95% CL upper limits on the production cross section times branching ratio for a $Z' \to t\bar{t}$ resonance with a relative decay width ($\Gamma_{Z'}/M_{Z'}$) of 10%. Limits are shown for the semileptonic analysis [2] (left) and the all-hadronic analysis [3] (right) as a function of the signal mass hypothesis.

Table 1: Observed mass exclusion range for the $X \to t\bar{t}$ signal models considered in the semileptonic and all-hadronic analyses. The corresponding values from the combination of $t\bar{t}$ searches in CMS at $\sqrt{s} = 8$ TeV [5] are also shown.

Observed mass exclusion region [TeV]

channel		Z′ (1% width)	Z′ (10% width)	Z′ (30% width)	KK gluon
ℓ + jets	(13 TeV) [2]	0.6 – 2.3	0.5 – 3.4	1.0 – 4.0	0.5 – 2.9
all-hadronic	(13 TeV) [3]	1.4 – 1.6	1.0 – 3.3	1.0 – 3.8	1.0 – 2.4
combination	(8 TeV) [5]	0.5 – 2.4	0.5 – 2.9	not tested	0.5 – 2.8

ACKNOWLEDGEMENTS

The financial support of the Spanish Ministerio de Economía y Competitividad, under project number FPA2014-53938-C3-3-R, is acknowledged.

References

[1] CMS Collaboration, JINST **3** (2008) S08004

[2] CMS Collaboration, CMS-PAS-B2G-15-002 [cdsweb.cern.ch/record/2138345]

[3] CMS Collaboration, CMS-PAS-B2G-15-003 [cdsweb.cern.ch/record/2160237]

[4] CMS Collaboration, CMS-PAS-JME-15-002 [cdsweb.cern.ch/record/2126325]

[5] CMS Collaboration, Phys. Rev. D **93** (2016) 012001